



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

January 12, 2017

Ms. Keshia Caswell  
Senior Regulatory Specialist for,  
Stepan Company  
22 West Frontage Road  
Northfield, IL 60093

Subject: PRIA Label Amendment – To add Influenza A Virus to the product labeling  
Product Name: BTC 2125M 10% Solution  
EPA Registration Number: 1839-86  
Application Date: May 24, 2016  
Decision Number: 520923

Dear Ms. Caswell:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Page 2 of 2  
EPA Reg. No. 1839-86  
Decision No. 520923

Your release for shipment of the product constitutes acceptance of these conditions.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. If you have any questions, please contact Karen M. Leavy by phone at (703)-308-6237, or via email at [Leavy.Karen@epa.gov](mailto:Leavy.Karen@epa.gov).

Sincerely,



Eric Miederhoff  
Product Manager 31  
Regulatory Management Branch I  
Antimicrobials Division (7510P)  
Office of Pesticide Programs

Enclosure


# BTC® 2125M 10% Solution

## FRONT PANEL

### ACTIVE INGREDIENTS

Alkyl (60% C <sub>14</sub> , 30% C <sub>16</sub> , 5% C <sub>12</sub> , 5% C <sub>18</sub> ) dimethyl benzyl ammonium chlorides.....	5.0%
Alkyl (68% C <sub>12</sub> , 32% C <sub>14</sub> ) dimethyl ethylbenzyl ammonium chlorides.....	5.0%
<b>(INERT)(OTHER) INGREDIENTS.....</b>	<b>90.0%</b>
<b>TOTAL.....</b>	<b>100.0%</b>

**NET CONTENTS(:)** XXX (oz)(ounces)(gallon(s))(gal(s))  
**(NET CONTENTS:** XX (oz)(ounce(s)) per packet)



## KEEP OUT OF REACH OF CHILDREN **DANGER**

FIRST AID	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
<b>If in Eyes:</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>If on Skin or Clothing:</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>• Call poison control center or doctor for treatment advice.</li></ul>
<b>If Swallowed:</b>	<ul style="list-style-type: none"><li>• Call poison control center or doctor immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.</li></ul>
<b>If Inhaled:</b>	<ul style="list-style-type: none"><li>• Move person to fresh air.</li><li>• If person is not breathing, call 911 or an ambulance then give artificial respiration, preferably by mouth-to-mouth, if possible.</li><li>• Call a poison control center or doctor for further treatment advice.</li></ul>
<b>(HOTLINE NUMBER</b> You may contact (xxx) xxx-xxxx for emergency medical treatment information.)	

### NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage.

*(Note to reviewer: Only one of the following statements is required. Referral to "Directions for Use" is required when directions for use are only found in booklet. Choose applicable statement noted in curly bracket.)*

See (side) (back)(s) panel for additional precautionary statements.

or

See booklet for {(complete) (additional)} information (and) (including) (Directions for Use)

## FRONT, BACK OR SIDE PANEL

EPA REG. NO. 1839-86  
EPA EST. NO. 1839-IL-01

### STEPAN COMPANY

22 West Frontage Road  
Northfield, IL 60093

Page 1 of 22  
Amend 8/22/16  
EPA Reg. No. 1839-86

Bracketed information is denoted as: { insert as noted }, ( optional text ), [ required qualifier ], <note>.  
The term "this product" used throughout this document may be replaced with the marketed product brand name.  
The term "x" is a place holder for numerical digits.  
Punctuation and plural/singular word forms may be adjusted to allow for grammatical correctness.  
Appropriate metric or imperial unit conversion may be added as optional supplemental information.

## **MARKETING CLAIMS**

### **General**

- DISINFECTANT
- DEODORIZER
- VIRUCIDE
- SANITIZER
- MILDEWSTAT (on hard non-porous, inanimate surfaces)
- CITRUS CANCKER CONTROL (*Not for Use in California*)
- Effective in the presence of 5% serum contamination
- For Home, Hospital, Institutional and Industrial Use
- For Farm, School, Dairy, Restaurant, Food Handling and Processing Areas, Equine, Poultry/Turkey Farm, Bar, Tavern and Institutional Kitchen Use
- Formulated for Effective Farm Premise Sanitation
- Formulated for Effective Poultry Premise Sanitation
- Formulated for Effective Swine Premise Sanitation
- Formulated for Effective Mushroom Farm Sanitation
- Formulated for Effective Veterinary Practice/Animal Care/Animal Laboratory Disinfection

<b>Front Panel Claim</b>	<b>Corresponding (Back)(Side) Panel Claim</b>
Kills 99.999% of bacteria in 60 seconds	(This product) (product name) is an effective sanitizer for use on food contact surfaces in 60 seconds at 200 ppm active quaternary against Escherichia coli, Escherichia coli O157 H7, Staphylococcus aureus, Yersinia enterocolitica, Listeria monocytogenes, Salmonella typhi, Shigella sonnei (ATCC 11060), Vibrio cholera, Methicilhn resistant Staphylococcus aureus (MRSA), Vancomycin resistant Enterococcus faecalis (VRE), Campylobacter jejuni and Klebsiella pneumoniae.
Formulated for third sink <sup>††</sup>	<sup>††</sup> Formulated for 3rd sink food contact surface sanitization
Make 512 gallons <sup>#</sup>	<sup>#</sup> 1 gallon makes 512 gallons of food contact surface sanitizer solution

- BTC<sup>®</sup> 2125M 10% Solution is designed for use in hospitals, nursing homes, hotels, schools, homes, athletic/recreational facilities/tanning spas, dairies, food processing plants, food service establishments, restaurants, bars, animal quarters and kennels where disinfection, sanitization and deodorization is of prime importance. When used as directed, this product is formulated to disinfect hard, non-porous, inanimate environmental surfaces: floors, walls, metal surfaces, stainless steel surfaces, glazed porcelain, glazed ceramic tile, plastic surfaces, bathrooms, shower stalls, bathtubs, cabinets, tables, chairs, tanning beds and telephones.
- This product contains no phosphorus.
- This product is for use in kitchens, bathrooms, and other household areas.
- This product has been formulated to aid in the reduction of cross-contamination on treated surfaces not only in hospitals, but in schools, institutions and industry.
- This product is an economical concentrate to be used with a mop and bucket, sponge, cloth, low pressure coarse sprayer, hand pump trigger sprayer or by soaking.
- This product may be applied through automatic washing systems, immersion tanks, foaming apparatus, low pressure sprayers, and fogging (wet misting) systems.

**BTC<sup>®</sup> 2125M 10% Solution** (product name) for use in:

- Hospitals, nursing homes, medical and dental offices and clinics, physician offices, operating rooms, isolation wards, and medical research facilities
- Patient care rooms and facilities, recovery, anesthesia, emergency rooms, X-ray cat labs, newborn nurseries, orthopedics, whirlpools, respiratory therapy
- EMS & fire facilities
- Day care centers and nurseries, sick rooms

- Acute care institutions, alternate care institutions, home healthcare institutions
- Life care retirement communities
- Restaurant, cafeterias, pizzerias, fast food operations, bars, taverns, cocktail lounges, food storage areas, food establishments, coffee shops, donut shops, bagel stores, liquor stores
- Supermarkets, convenience stores, retail and wholesale establishments, department stores, shopping malls, gift shops, video stores, bookstores, dressing rooms, laundries, photo copy centers, bicycle shops, auto repair centers, toy stores
- Computer manufacturing sites, toy manufacturing sites
- Crime scenes, funeral homes, mortuaries
- Police stations, courthouses, municipal government buildings
- Bus stations, train stations, transportation terminals, airports, heliports
- Institutional facilities, laboratories, factories
- Lodging establishments, hotels, motels, bed & breakfast, dormitories
- Public restrooms, travel rest areas, shower rooms, shower stalls, bathrooms
- Kitchens, bathrooms and other household areas
- Schools, colleges, universities, classrooms, education institutions, school administration offices
- Athletic facilities, sports stadiums, convention centers, amphitheaters, health clubs, spas, tanning salons/spas, tanning beds, massage/facial salons, hair/nail/pedicure salons, barber/beauty shops, salons, tattoo parlors
- Museums, art galleries, post offices, performance/theater centers, banks, libraries, movie theaters, theaters
- Recycling centers and equipment
- Camp grounds, play grounds, recreational facilities, picnics facilities, sports arenas, sports complexes, sports stadiums
- Food processing plants, dairy farms, hog farms, equine farms, poultry & turkey farms, egg processing plants, meat/poultry processing plants, meat/poultry producing establishments, mushroom farms
- Fishery, milk, citrus, ice cream, potato processing plants
- Veterinary clinics, animal life science laboratories, kennels, dog/cat animal kennels, breeding and grooming establishments, pet animal quarters, zoos, pet shops, tack shops and other animal care facilities
- Household and automotive garages, boats, ships, campers, trailers, mobile homes, cars, trucks, buses, trains, taxis and airplanes
- Commercial florist and flower shops

This product may be used on washable hard, non-porous surfaces:

- Food preparation and storage areas
- Dishes, glassware, silverware, cooking utensils, plastic and other non-porous cutting boards, coolers, ice chests
- Kitchen equipment: food processors, blenders, cutlery, utensils
- Countertops, Slurppy<sup>®</sup> Machines, stovetops, sinks, appliances, refrigerators, ice machines
- Glass, metal, stainless steel, glazed porcelain, glazed ceramic, sealed fiberglass, sealed granite, sealed marble, plastic, chrome, vinyl
- Enameled surfaces, painted woodwork, Formica<sup>®</sup>, vinyl and plastic upholstery
- Tables, chairs, desks, bed frames, lifts, washable walls, cabinets, doorknobs, garbage cans
- Exhaust fans, refrigerated storage equipment, display equipment
- Large inflatable non-porous plastic and rubber structures such as animals, promotional items, moonwalk, slides, obstacle course play and exercise equipment
- Beer fermentation and holding tanks, bottling or pre-mix dispensing equipment
- Citrus processing equipment and holding tanks
- Picnic tables (non-wooden) and outdoor furniture (except cushions and wood frames)
- Telephones
- High chairs
- Drinking fountains
- Shower stalls, shower doors and curtains, bathtubs, glazed ceramic tile, glazed porcelain, chrome plated fixtures, toilets, toilet bowl surfaces, urinals, portable toilets, porcelain tile, restroom fixtures, bathroom fixtures,

- metal surfaces, stainless steel surfaces
- Kennels, kennel runs, cage floors, cages

## **DISINFECTION CLAIMS**

- Kills germs\*\*
- (Kills)(disinfects)(against){insert pathogen from DISINFECTION heading in chart on page 7-8}
- When used as directed, this product is a concentrated Hospital Use disinfectant that is an effective broad spectrum bactericide and virucide\*.
- This product is a versatile Disinfectant for Veterinarian, Animal Care and Agricultural premises.
- This product is a complete, chemically balanced disinfectant/sanitizer that provides clear use- solutions even when diluted in hard water.
- This product is for use as a hard, non-porous surface hospital disinfectant at 547 ppm active quaternary and as a general disinfectant at 469 ppm active quaternary.
- This product is a versatile broad spectrum disinfectant formulated for use in Ultrasonic Baths (Ultrasonic cleaning units).
- This product is a versatile cleaner and broad spectrum disinfectant formulated for use on bath and therapy equipment (Whirlpools).
- This product is an effective antimicrobial cleaner designed for use by wholesale and retail florists, shippers and greenhouses.
- When used as directed, this product will disinfect hard, non-porous surfaces such as flower buckets, floors, walls of coolers, design and packing benches and countertops.
- Use this product to clean, disinfect and deodorize flower buckets, walls and floors of coolers, shippers, greenhouse packing areas, garbage pails and other areas where obnoxious odors develop.
- This product may be used as a general purpose antimicrobial detergent in florist shops, wholesale florists, shippers, greenhouse packing areas and other commercial floriculture places for efficient cleaning and antimicrobial action against certain bacteria which cause:
  - Plugging of stems with slime, which reduces uptake of water for various sensitive flowers including roses, chrysanthemums, gladioli and tulips.
  - Production of ethylene gas, which may injure blooms of the various sensitive flowers including carnations, snapdragons, some orchids, babies breath, sweet peas, freesia and alstroemena.
- This product is recommended for Poultry Premise Disinfection (Hatcheries), Egg Receiving Areas, Egg Holding Areas, Setter Room, Tray Dumping Area, Trays, Buggies, Racks, Egg Flats, Chick Holding room, Poultry Buildings, Ceilings, Sidewalls and Floors, Drinkers, and all other Poultry House related Equipment, and all other hard, non-porous surfaces in the Hatchery Environment.
- Use this product for Swine Premise Disinfection: Waterers and Feeders, Hauling Equipment, Dressing Plants, Loading Equipment, Farrowing Barns and Areas, Nursery, Blocks, Creep Area, Chutes.
- This product meets AOAC Use Dilution Test Standards for hospital disinfectants.
- Respiratory illnesses attributable to Pandemic 2009 H1N1 are caused by Influenza A virus. This product is a broad-spectrum hard surface disinfectant that has been shown to be effective against (Influenza A virus tested and listed on the label) and is expected to inactivate all Influenza A viruses including Pandemic 2009 H1N1 (formerly called swine flu).
- This product has demonstrated effectiveness against Influenza A virus and is expected to inactivate all Influenza A viruses including Pandemic 2009 H1N1 Influenza A virus.
- This product has demonstrated effectiveness against (Influenza A virus tested and listed on the label) and is expected to inactivate all Influenza A viruses including Pandemic 2009 H1N1 (formerly called swine flu).
- Kills Pandemic 2009 H1N1 Influenza A virus (formerly called swine flu).
- Kills Pandemic 2009 H1N1 Influenza A virus.
- At 1 ounce per 4 gallons, this sanitizer fulfills the criteria of Appendix F of the Grade A Pasteurized Milk Ordinance 1978 Recommendations of the U. S. Public Health Services in waters up to 650 ppm of hardness calculated as CaCO<sub>3</sub> when evaluated by the AOAC Germicidal and Detergent Sanitizer Method against Escherichia coli and Staphylococcus aureus.
- **KILLS HIV-1, HBV and HCV ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care settings (hospitals, nursing homes) or other settings in which there is an expected likelihood of soiling of

inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS), Hepatitis B virus (HBV) and Hepatitis C virus (HCV).

- **EFFICACY TESTS HAVE DEMONSTRATED THAT THIS PRODUCT IS AN EFFECTIVE BACTERICIDE AND VIRUCIDE IN THE PRESENCE OF ORGANIC SOIL (5% BLOOD SERUM).**

## **SANITIZATION CLAIMS**

- (An effective sanitizer)(sanitizes) against {insert pathogen from SANITIZATION heading in chart on page 7-8}
- This product is for use for the sanitization of shell eggs intended for food in shell egg and egg product processing plants when used as directed.
- This product is for use as a Hatching-Egg sanitizer, with best results achieved in water temperatures ranging from 78°-110°F.
- This product is for use as a sanitizer on dishes, glassware, utensils and dairy processing equipment at 200-400 ppm active quaternary without potable water rinse.
- This product is for use as a sanitizer in public eating establishments, dairies and on food processing equipment at 200-400 ppm active quaternary without potable water rinse.
- This product is for use as a sanitizer on bottling and beverage dispensing equipment. This product is for use as a sanitizer in sanitary filling of bottles and cans.
- This product is for use as a sanitizer for bottles or cans in the final rinse application and for external spraying of filling and closing machines.
- This product is for use as a sanitizer on beer fermentation and holding tanks.
- Use this product to sanitize and disinfect non-porous salon/barber tools and instruments: combs, brushes, scissors, blades, and manicure instruments.
- This product meets AOAC Germicidal & Detergent standards for sanitizing previously cleaned food-contact surfaces.
- This product is an effective sanitizer for use on food contact surfaces in 60 seconds at 200-400 ppm active quaternary against Staphylococcus aureus, Escherichia coli, Escherichia coli O157:H7, Yersinia enterocolitica, Listeria monocytogenes, Salmonella typhi, Shigella sonnei, Vibrio cholera, Methicillin resistant Staphylococcus aureus (MRSA), Vancomycin resistant Enterococcus faecalis (VRE), Campylobacter jejuni and Klebsiella pneumoniae.
- Regular, effective cleaning and sanitization of equipment, utensils and work or dining surfaces that could harbor hazardous microorganisms minimizes the probability of contaminating food during preparation, storage or service. Effective cleaning will remove soil to prevent the accumulation of food residues, which may decompose or support the rapid development of food poisoning organisms or toxins. Application of effective sanitization procedures reduces the number of those disease causing organisms that may be present on equipment and utensils after cleaning. Effective sanitization reduces the potential for the transfer of disease, either directly through tableware such as glasses, cups and flatware or indirectly through food.
- To prevent cross-contamination, kitchenware and equipment food contact surfaces must be washed, rinsed with potable water and sanitized after each use and following any interruption of operation during which time contamination may have occurred.
- Where equipment and utensils are used for the preparation of foods on a continuous production-line basis, utensils and equipment food-contact surfaces must be washed, rinsed with potable water and sanitized at intervals throughout the day on a schedule based on food temperature, type of food, and the amount of food particle accumulation.
- Sanitization of public eating establishment and dairy food contact surfaces is regulated under 40CFR180.940.

## **FUNGI CLAIMS**

- Use this product on multi-touch surfaces responsible for cross-contamination. This product is effective at controlling mold and mildew odor on shower curtains.

## **DEODORIZATION & CLEANING CLAIMS**

- This product deodorizes those areas, which generally are hard to keep fresh smelling, such as garbage storage areas, empty garbage bins and cans, pet areas and any other areas, which are prone to odors caused by microorganisms.
- This product will not leave a grit or soap scum.
- This product may be relied on to deodorize coolers, buckets, garbage pails and other areas where obnoxious odors may develop.
- This product is for use in work areas such as tool rooms and garages for odor control and light duty cleaning.
- This product provides long lasting freshness against tough (pet) odors from litter boxes and pet accidents.
- This product neutralizes on contact musty odors and tough odors from smoke, pet accidents, and spills.
- This product is formulated to effectively eliminate offensive odors caused by mold and mildew.
- This product (controls) (reduces) (eliminates) (destroys) odors to make your home (kitchen) (bath- room) sanitary.
- This product is effective against household odors by animal waste, septic tank or sewage backup, smoke and bathroom and kitchen odors.

## **CLAIM QUALIFIERS**

\*\*Kills Staphylococcus aureus, Salmonella enterica, Listeria monocytogenes, Yersinia enterocolitica, Pseudomonas aeruginosa, Vancomycin intermediate resistant Staphylococcus aureus (VISA), Influenza A<sub>2</sub>/Japan, Herpes Simplex Type 1, Adenovirus Type 5, Vaccinia Virus, Avian Influenza A/Turkey/Wisconsin, Human Immunodeficiency Virus Type 1 (HIV-1), Newcastle Disease Virus, Laryngotracheitis Virus, Hepatitis B Virus (HBV), Hepatitis C Virus (HCV), Bovine Viral Diarrhea Virus (BVDV), SARS associated Coronavirus, Porcine Respiratory & Reproductive Syndrome Virus (PRRSV) and Human Coronavirus.

*<For labels that list medical premises and metal and/or stainless steel surfaces, one of the following statements must be used:>*

This product is not to be used as a terminal sterilant/high-level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body.

OR

This product is not for use on critical and semi-critical medical device surfaces.

## **MIRCO-ORGANISMS**

This product has been tested and found to be efficacious against the following micro-organisms on hard, non-porous surfaces:

<b>DISINFECTION</b>		
<b>Bactericidal Activity (Bacteria)</b>	<b>ATCC No</b>	<b>Contact Time</b>
Pseudomonas aeruginosa	(ATCC 15442)	10 minutes
Staphylococcus aureus	(ATCC 6538)	10 minutes
Salmonella enterica	(ATCC 10708)	10 minutes
Listeria monocytogenes	(ATCC 35152)	10 minutes
Vancomycin intermediate resistant Staphylococcus aureus (VISA)		10 minutes
Yersinia enterocolitica	(ATCC 23715)	10 minutes
<b>Virucidal Activity (Viruses)</b>	<b>ATCC No</b>	<b>Contact Time</b>
Adenovirus Type 5		10 minutes
Avian influenza A/Turkey/Wisconsin	(ATCC VR-798)	10 minutes
Influenza A Virus (H7N9)		10 minutes
Bovine Viral Diarrhea Virus (BVDV)		10 minutes
Hepatitis B Virus (HBV)		10 minutes
Hepatitis C Virus (HCV)		10 minutes
Herpes Simplex Type 1		10 minutes
Human Coronavirus (Strain 229E)	(ATCC VR-740)	10 minutes
Human Immunodeficiency Virus (HIV-1)(Strain HTLV-IIIB)		10 minutes
Human Influenza		10 minutes
Influenza A2 (Japan 305/57)		10 minutes
Pandemic 2009 H1N1 Influenza A Virus		10 minutes
Laryngotracheitis Virus (LT-IVAX)		10 minutes
SARS Associated Coronavirus (ZeptoMetrix)		10 minutes
Vaccinia Virus		10 minutes
Newcastle disease (strain H. J. Roakin, 1946)		10 minutes
Porcine Respiratory & Reproductive Syndrome Virus (PRRSV) (Strain NVSL)		10 minutes
<b>FOOD CONTACT SANITIZATION</b>		
Escherichia coli	(ATCC 11229)	60 seconds
Escherichia coli O157:H7	(ATCC 43895)	60 seconds
Staphylococcus aureus	(ATCC 6538)	60 seconds
Yersinia enterocolitica	(ATCC 23715)	60 seconds
Listeria monocytogenes	(ATCC 35152)	60 seconds
Methicillin resistant Staphylococcus aureus (MRSA)	(ATCC 33592)	60 seconds
Vancomycin resistant Enterococcus faecalis (VRE)	(ATCC 51299)	60 seconds
Campylobacter jejuni	(ATCC 29428)	60 seconds
Klebsiella pneumoniae	(ATCC 4352)	60 seconds
Salmonella typhi	(ATCC 6539)	60 seconds
Shigella sonnei	(ATCC 11060)	60 seconds

Bracketed information is denoted as: { insert as noted }, ( optional text ), [ required qualifier ], <note>.  
 The term "this product" used throughout this document may be replaced with the marketed product brand name.  
 The term "x" is a place holder for numerical digits.  
 Punctuation and plural/singular word forms may be adjusted to allow for grammatical correctness.  
 Appropriate metric or imperial unit conversion may be added as optional supplemental information.

Vibrio cholera	(ATCC 14035)	60 seconds
<b>NON-FOOD CONTACT SANITIZATION</b>		
Klebsiella pneumoniae	(ATCC 4352)	30 seconds
Staphylococcus aureus	(ATCC 6538)	30 seconds
Listeria monocytogenes	(ATCC 35152)	30 seconds

## **DIRECTIONS FOR USE**

### **DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**GENERAL DEODORIZATION:** To deodorize, add 3 ounces of (this product or product name) to 5 gallons of water (or equivalent use dilution) (469 ppm active quat). Let air dry.

#### **FOOD CONTACT SURFACE SANITIZATION DIRECTIONS:**

**Sanitization:** Recommended for use in restaurants, dairies, food processing plants, and bars.

When used as directed this product is an effective sanitizer at an active quaternary concentration of 200-400 ppm when diluted in water up to 650 ppm hardness (CaCO<sub>3</sub>) in public eating establishments, dairies and food processing plants against Escherichia coli (ATCC 11229), Escherichia coli O157:H7 (ATCC 43895), Staphylococcus aureus (ATCC 6538), Yersinia enterocolitica (ATCC 23715), Listeria monocytogenes (ATCC 35152), Salmonella typhi (ATCC 6539), Shigella sonnei (ATCC 11060), Vibrio cholera (ATCC 14035), Methicillin resistant Staphylococcus aureus (MRSA) (ATCC 33592), Vancomycin resistant Enterococcus faecalis (VRE) (ATCC 51299), and Campylobacter jejuni (ATCC 29428).

*<NOTE: If Klebsiella pneumoniae is included, the following paragraph must be included.>*

This product is an effective sanitizer against Klebsiella pneumoniae (ATCC 4352) at a 200 ppm active quaternary concentration when diluted in water up to 500 ppm hardness (CaCO<sub>3</sub>). At an active quaternary concentration of 300 ppm, it is an effective sanitizer against Klebsiella pneumoniae (ATCC 4352) when diluted in water up to 650 ppm hardness (CaCO<sub>3</sub>).

Remove all gross food particles and soil from areas, which are to be sanitized, with a good detergent, pre-flush, pre-soak or pre-scrape treatment. Rinse with a potable water rinse.

Prepare a fresh solution daily or more frequently as soil is apparent.

#### **NON-FOOD CONTACT SURFACE SANITIZATION DIRECTIONS:**

This product is an effective sanitizer for pre-cleaned, hard, inanimate, non-porous surfaces: walls, floors, bathroom surfaces, shower stalls, shower doors, shower curtains, cabinets, toilets, chrome plated fixtures, urinals, doorknobs, garbage cans, exercise equipment, wrestling mats, glass surfaces, stainless steel surfaces, metal surfaces, glazed ceramic tile, glazed fiberglass, drains, telephones, telephone keypads, animal cages, cages, kennel runs, automotive garages, boats, ships, automobile interior surfaces: steering wheels, seat belt buckles and housings, non-porous surfaces of child restraint seats, door handles, non-porous surfaces of visors, mirrors, dashboards, gear shift levers, accessory control knobs, vinyl car seats and head restraints. A sanitizer use-solution of (0.25 oz. per gallon of water)(1 oz. per 4 gallons of water)(2.5 ozs. per 10 gallons of water)(5 ozs. per 20 gallons of water) (200 ppm active quaternary) will kill (99.9%) of Klebsiella pneumoniae (ATCC 4352), Staphylococcus aureus (ATCC 6538) and Listeria monocytogenes (ATCC 35152).

Sanitization will occur only on pre-cleaned surfaces. The sanitizer use-solution can be applied to pre-cleaned, hard, inanimate, non-porous surfaces with a cloth, sponge, mop, sprayer or by flooding or immersion.

For sprayer applications, use a coarse (mist) pump or trigger sprayer. Spray 6-9 inches from the surface. Rub with brush, sponge or cloth. Do not breathe spray (mist).

**NOTE:** With spray applications, cover or remove all food products.

Treated surfaces must remain wet for 30 seconds (1/2 minute) and (wiped with a sponge) (wiped with a mop) (wiped with a cloth) (and/or) (allowed to air dry).

**DISINFECTION** - To disinfect inanimate, hard non-porous surfaces apply use-solution with mop, cloth, sponge, low pressure coarse sprayer or hand pump trigger sprayer so as to wet all surfaces thoroughly. Allow to remain wet for 10 minutes, then remove excess liquid. For heavily soiled areas, a pre-cleaning step is required. Prepare a fresh solution for each use. Surfaces which may contact food must be rinsed thoroughly with potable water after use.

For low pressure coarse sprayer and hand pump trigger sprayer applications, spray 6-8 inches from the surface; rub with brush, sponge or cloth. Do not breathe spray.

NOTE: For spray applications, cover or remove all food products.

**GENERAL DISINFECTION** - Add 3 ounces of this product per 5 gallons of water (or equivalent dilution) for disinfection against Staphylococcus aureus, Salmonella enterica, Listeria monocytogenes, and Yersinia enterocolitica.

**General Disinfectant/Non-Medical Dilution Chart**

Product Amount (ounces)	Water Volume
0.15	1.0 quart
0.3	0.5 gallon
0.6	1.0 gallon
1.5	2.5 gallons
3.0	5.0 gallons
6.0	10.0 gallons
x	x gallons

**HOSPITAL DISINFECTION** - Add 3.5 ounces of (this product or product name) per 5 gallons of water (or equivalent dilution) for disinfection against Pseudomonas aeruginosa and Vancomycin intermediate resistant Staphylococcus aureus (VISA), as well as the organisms listed under the GENERAL DISINFECTION section.

**Hospitals, Dental Offices, Nursing Homes and other Health Care Institutions Disinfection and Virucide Dilution Chart**

Product Amount (ounces)	Water Volume
0.175	1.0 quart
0.35	0.5 gallon
0.7	1.0 gallon
1.75	2.5 gallons
3.5	5.0 gallons
7.0	10.0 gallons
x	x gallons

Bracketed information is denoted as: { insert as noted }, ( optional text ), [ required qualifier ], <note>.  
 The term "this product" used throughout this document may be replaced with the marketed product brand name.  
 The term "x" is a place holder for numerical digits.  
 Punctuation and plural/singular word forms may be adjusted to allow for grammatical correctness.  
 Appropriate metric or imperial unit conversion may be added as optional supplemental information.

**\*VIRUCIDAL ACTIVITY** - This product, when used on environmental, inanimate, hard non-porous surfaces at 3.5 ounces per 5 gallons (or equivalent dilution) of water exhibits effective virucidal activity against Influenza A<sub>2</sub>/Japan (representative of the common flu virus), Herpes Simplex Type 1 (causative agent of fever blisters), Adenovirus Type 5 (causative agent of upper respiratory infections), Vaccinia Virus (representative of the pox virus group), Avian Influenza A/Turkey/Wisconsin (causative agent of an acute avian lower respiratory tract infection), HIV-1 (Human Immunodeficiency Virus Type 1), Newcastle Disease Virus, Laryngotracheitis Virus, Hepatitis B Virus (HBV), Hepatitis C Virus (HCV), Bovine Viral Diarrhea Virus (BVDV), Human Coronavirus (ATCC VR-740, Strain 229E), Porcine Respiratory & Reproductive Syndrome Virus (PRRSV) (Strain NVSL) and SARS associated Coronavirus.

*<required if making an HIV, HBV, and/or HCV claim on the product label>*

**SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HBV and HCV OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.**

**PERSONAL PROTECTION:** When handling items soiled with blood or body fluids use disposable latex gloves, gowns, masks, and eye coverings.

**CLEANING PROCEDURES:** Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this product.

**CONTACT TIME:** Allow surface to remain wet for 10 minutes.

**DISPOSAL OF INFECTIOUS MATERIALS:** Blood and other body fluids must be autoclaved and disposed of according to local regulations for infectious waste disposal.

**MILDEWSTAT:** To control mold and mildew on pre-cleaned, hard, non-porous surfaces add 3 ounces of this product per 5 gallons of water. Apply solution with a cloth, mop, sponge, or hand pump trigger sprayer making sure to wet all surfaces completely. Let air dry. Prepare a fresh solution for each use. Repeat application at weekly intervals or when mildew growth appears.

*<The following language is required for products supplied with supplemental labeling: inserts, hang tags and reference sheets.>*

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Labeling must be in the possession of the user at the time of using the product. See container label for FIRST AID and PRECAUTIONARY STATEMENTS and complete DIRECTIONS for USE before applying. Use of this product is subject to the precautions and limitations imposed by the label affixed to the container for this product.

*<The following language is required to appear on the container label:>*

See (insert) (hangtag) (reference sheet) for (complete) (additional) information.

*<Alternative language for bag in the box dispensing equipment>*

**Directions For Bag-In Box Containers.**

**HOW TO USE THIS PACKAGE:** This package is designed to be used with dilution control systems only. Open package and connect to dispense according to directions on the box.

**TRIGGER SPRAYERS:** Fill bottle from dispenser. Apply to surfaces according to directions above.

**MOP BUCKETS:** Fill bucket from dispenser. Set up "Wet Floor" signs. Mop floor surfaces as specified in directions above.

**SITE & APPLICATION SPECIFIC DIRECTIONS FOR USE**

**Non-acid toilet bowl and urinal disinfectant directions:** Remove heavy soil prior to disinfecting. Empty toilet bowl or urinal and apply a use-solution of 3.5 ounces per 5 gallons of water (or equivalent dilution) to exposed surfaces including under the rim with a cloth, mop, sponge or hand pump trigger sprayer. Brush or swab thoroughly, then allow to remain wet for 10 minutes and flush.

**To disinfect food processing premises:** floors, walls, and storage areas, add 3 ounces of this product per 5 gallons of water (or equivalent dilution). For heavily soiled areas, a pre-cleaning step is required. Apply solution with a mop, cloth, sponge or hand pump trigger sprayer so as to wet all surfaces thoroughly. Allow to remain wet for 10 minutes, then remove excess liquid. Before using this product, food products and

packaging materials must be removed from the area or carefully protected. After use, all surfaces in the area must be thoroughly rinsed with potable water.

**To disinfect (food service establishment) (or) (restaurant food contact surfaces):** floors, walls, storage areas, countertops, outside of appliances, tables, add 3 ounces of this product per 5 gallons of water (or equivalent dilution). For heavily soiled areas, a pre-cleaning step is required. Apply solution with a cloth, sponge or hand pump trigger sprayer so as to wet all surfaces thoroughly. Allow the surface to remain wet for 10 minutes, then remove excess liquid and rinse the surface with potable water. Do not use on utensils, dishes, glasses or cookware.

**Disinfecting potato storage area and equipment:** Remove all potatoes prior to disinfection of potato storage area or equipment. Pre-clean hard surfaces by removing heavy soil or gross filth. Follow general disinfection (3 oz. per 5 gal.) procedure described above. All treated surfaces must be thoroughly rinsed with potable water prior to reuse.

**For use as a general disinfectant on brewery premises (non-food contact surfaces):** Floors, walls, and storage areas add 3 ounces of (this product or product name) per 5 gallons of water (or equivalent dilution). For heavily soiled areas, a pre-cleaning step is required. Apply use-solution with a mop, cloth, sponge, low pressure coarse sprayer or hand pump trigger sprayer so as to wet all surfaces thoroughly. Allow to remain wet for 10 minutes, then remove excess liquid. Before using product, food products (beer, wort, hops, mash, grain, brew ingredients) must be removed from the area or carefully protected. After use, all surfaces must be thoroughly rinsed with potable water.

**Ultrasonic bath disinfectant directions:** Use (this product or product name) to disinfect hard, non-porous, non-critical instruments/objects compatible with Ultrasonic cleaning units. Pour fresh use-solution of 3.5 ounces per gallon of water (or equivalent dilution) directly into bath chamber. Pre-clean soiled objects. Place objects into unit and operate for a minimum of 10 minutes, according to manufacturer's use directions. Remove objects and rinse with sterile water (sterile water for injection), or allow to air dry. Prepare a fresh solution for each use.

**Whirlpool bath disinfection directions:** After using whirlpool unit, drain and fill with a use-solution of 3.5 ounces of (this product or product name) per 5 gallons (or equivalent dilution) to just cover the intake valve. Start pump to circulate solution. Wash down the unit sides, seat of the chair lift, and any related equipment with a clean swab, brush or sponge. Treated surfaces must remain wet for 10 minutes. After the unit has been thoroughly disinfected, drain the solution from the unit and rinse disinfected surfaces with fresh water. Repeat for heavily soiled units.

**Commercial Florist Use Directions:** To clean, disinfect and deodorize hard non-porous surfaces in one step, prepare use-solution by adding (mixing) 3.5 ounces per 5 gallons of water (or equivalent dilution). For heavy-duty use, add (mix) 4 ounces per gallon of water.

Remove all leaves, petals, garbage and refuse. Pre-clean surfaces using pressurized water where possible. Apply use-solution to hard (inanimate) non-porous surfaces thoroughly wetting surfaces as recommended and required, with a cloth, mop, brush, sponge, hand pump trigger sprayer or low pressure coarse sprayer.

For heavily soiled areas, a preliminary cleaning is required.

For sprayer applications, spray 6-8 inches from surface; rub with brush, sponge or cloth. Do not breathe spray mist.

Treated surfaces must remain wet for 10 minutes. Allow to air dry.

Prepare a fresh solution at least daily or sooner if use-solution becomes visibly dirty.

**Disinfection of poultry/turkey equipment, swine quarters, animal quarters and kennels:** For disinfection of pre-cleaned poultry equipment (brooders, watering founts, feeding equipment), animal quarters and kennels, apply a use-solution of 3.5 ounces of (this product or product name) per 5 gallons of water (or equivalent dilution). Remove all poultry, animals, and feed from premises, trucks, coops, and crates. Remove all litter and droppings from floors, walls and surfaces of facilities occupied or traversed by poultry or animals. Empty all troughs, racks, and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate the surfaces with the disinfecting solution for a period of 10 minutes. Immerse all halters, ropes, and other types of equipment used in handling and restraining animals as well as forks, shovels, and scrapers used for removing litter and

manure. Ventilate building, coops, and other closed spaces. Do not house poultry or animals or employ equipment until treatment has been absorbed, set or dried. All treated equipment that will contact feed or drinking water must be rinsed with potable water before reuse.

**Hatcheries:** To disinfect and provide virucidal\* activity, use 3.5 ounces of (this product or product name) per 5 gallons of water (or equivalent dilution). Apply to hatcheries, setters, trays, racks, carts, sexing tables, delivery trucks and other hard, non-porous surfaces. Leave all treated surfaces exposed to the use-solution for a minimum of 10 minutes and allow to air dry. For heavily soiled areas, a pre-cleaning step is required.

**Vehicles:** Clean all vehicles including mats, crates, cabs and wheels with high pressure water and (this product or product name). For spray applications use a coarse spray device. Use 3.5 ounces of (this product or product name) per 5 gallons of water (or equivalent dilution) for disinfection and virucidal\* activity. Leave all treated surfaces exposed to the use-solution for a minimum of 10 minutes and allow to air dry.

**Dressing plant disinfectant/virucidal\* directions:** Apply a use-solution of 3.5 ounces per 5 gallons of water (or equivalent dilution) to equipment, utensils, walls and floors in poultry and animal dressing plants. Equipment and utensils must remain wet for a minimum of 10 minutes then thoroughly rinsed with potable water before resuming operations. Disinfect all rooms, exterior walls and loading platforms of dressing plants. For heavily soiled areas, a pre-cleaning step is required.

**Farm premise disinfection:** For use in Equine, Dairy and Hog Farms.

1. Remove all animals and feed from premises, vehicles and enclosures.
2. Remove all litter and manure from floors, walls and surfaces of barn, pens, chutes and other facilities and fixtures occupied or traversed by animals.
3. Empty all troughs, racks and other feeding and watering appliances.
4. Thoroughly clean all surfaces with a compatible soap or detergent and rinse with potable water.
5. Saturate all hard, non-porous surfaces with a use-solution of 3.5 ounces per 5 gallons of water (or equivalent dilution) and allow to remain wet for a minimum of 10 minutes.
6. Immerse all halters, ropes and other types of equipment used in handling and restraining animals as well as forks, shovels, scrapers used in removing litter and manure.
7. Ventilate buildings, cars, trucks, boats and other closed spaces. Do not house livestock or employ equipment until treatment has been absorbed, set and dried.
8. Thoroughly scrub all treated feed racks, mangers, troughs, automatic feeders, fountains and waterers with a compatible soap or detergent and rinse with potable water before reuse.

**Shoe bath sanitizer directions:** To prevent tracking harmful organisms into animal areas, and the packaging and storage areas of food plants, shoe baths containing one inch of freshly made use-solution should be placed at all entrances to buildings, hatcheries and all entrances to the production and packaging rooms. Scrape waterproof shoes and place in a 0.25 ounce of (this product or product name) per gallon of water (or equivalent use dilution) (200 ppm active solution) for 60 seconds prior to entering area. Change use-solution in the bath daily or sooner if use-solution appears dirty.

**Shoe foam sanitizer directions:** To prevent tracking harmful organisms into animal areas, and the packaging and storage areas of food plants, apply a foam layer approximately 0.5 to 2 inches thick made from a solution of 1 to 1.5 ounces per gallon of water (or equivalent use dilution) (800 to 1200 ppm active quaternary) at all entrances to buildings, hatcheries, production and packaging rooms by using a foam generating machine or aerator to apply foam layer. Follow the foaming directions as specified by the manufacturer of the foam generator/aerator. Scrape waterproof shoes. Stand and/or walk through foamed area for 60 seconds prior to entering area. Foam area should be washed and replaced daily or when it appears dirty.

**Veterinary clinics/animal life science laboratory/pet shop/kennels/breeding and grooming establishments/tack shops disinfection directions:** For cleaning and disinfection of the following hard, non-porous surfaces: equipment not used for animal food or water, utensils, instruments, cages, kennels, stables and catteries. Remove all animals and feeds from premises, animal transportation vehicles, crates, etc. Remove all litter, droppings and manure from floors, walls and surfaces of facilities occupied or traversed by animals. Thoroughly clean all surfaces with a compatible soap or detergent and rinse with potable water. Saturate all hard, non-porous surfaces with a use-solution of 3.5 ounces of (this product or product name) per 5 gallons of water (or equivalent dilution) for 10 minutes minimum. Ventilate buildings

and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried. Scrub all treated feed racks, automatic feeders, waterers and other equipment that dispenses food or water with a compatible soap or detergent and thoroughly rinse with potable water before reuse.

**Salon (barber) instruments and tools disinfection (bactericide/virucide\*) directions:** Mix 3.5 ounces of (this product or product name) per 5 gallons of water (or equivalent dilution) to disinfect, hard, non-porous instruments and tools. Completely immerse combs, brushes, rollers, razors, scissors, blades, manicure and other salon instruments and tools for 10 minutes. For heavily soiled instruments and tools, a preliminary cleaning is required. Rinse thoroughly and dry before use. Prepare a fresh solution daily or more often if the solution becomes visibly cloudy or soiled.

**NOTE:** Plastics may remain immersed until ready to use. Stainless steel shears and instruments must be removed after 10 minutes, rinsed, dried and kept in a clean, non-contaminated receptacle. Prolonged soaking may cause damage to metal instruments.

**To sanitize barber/salon instruments and tools:** Immerse pre-cleaned barber/salon instruments and tools (combs, brushes, razors, scissors, blades and manicure instruments) in a use-solution of 1 ounce of (this product or product name) per 4 gallons of water (or equivalent dilution) for at least 60 seconds (one minute). Rinse thoroughly and dry before use. Prepare a fresh solution daily or more often if solution becomes visibly cloudy or soiled.

**NOTE:** Plastics may remain immersed until ready to use. Stainless steel shears and instruments must be removed after 10 minutes, rinsed, dried and kept in a clean, non-contaminated receptacle. Prolonged soaking may cause damage to metal instruments.

**To Control Mold and Mildew on Large, Inflatable, Non-Porous Plastic and Rubber Structures (animals, promotional items, moonwalk, slides, obstacle course play equipment, exercise equipment, and wrestling mats).** Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate surfaces with a use-solution of 3 ounces of (this product or product name) per 5 gallons of water (0.6 ounce per gallon of water) (469 ppm active quaternary) (or equivalent dilution) for a period of 10 minutes. Ventilate buildings and other closed spaces. Do not use equipment until treatment has been absorbed, set or dried.

**For Deodorizing Garbage Cans, Garbage Trucks, Industrial Waste Receptacles and Garbage Handling Equipment:** It is especially important to pre-clean for this product to perform properly. Apply a wetting concentration of 3.25 ounces of (this product or product name) per gallon of water (or equivalent use dilution) (2500 ppm active quaternary).

**For Odors Caused by Dogs, Cats and Other Domestic Animals:** Use on rugs, floors, walls, tile, cages, litter boxes, mats, floor coverings, or any surface soiled by a pet. Test a small inconspicuous area first. Blot problem area. Then follow directions for "General Deodorization".

**Cleansing of Body Surfaces and Body Orifices of Human Remains.** To cleanse away skin secretions and accompanying malodor, to insure the removal of all soil and bloodstains, and to remove and reduce surface contamination, apply 3.5 ounces of (this product or product name) to 5 gallons of water (or equivalent dilution) (547 ppm active quaternary) to the surfaces and body openings, natural or artificial. Allow 10 minute contact time for optimal results. Bathe the entire body using sponge or washcloth. A soft brush may be employed on surfaces other than the face. Prepare a fresh solution for application to each remains.

**Drain Cleaning:** For cleaning drains, dilute 1.25 ounces of (this product) (product name) per one gallon of water. Pour solution down drain being sure to coat all sides of drainpipe. Allow all treated surfaces exposed to solution to air dry.

**Cleansing Aids:** For cleaning plastic brooms, brushes, squeegees, wet/dry vacuums and condensate removal equipment, dilute 1.25 ounces of (this product) (product name) per one gallon of water to provide 1000 ppm active quaternary. The cleaning aids should either be stored dry or in a use-solution of (this product) (product name) of 1000 ppm active quaternary.

**Boot Cleaning:** Wash, foam or coarse spray boots with solution of 1.25 ounces of (this product) (product name) per one gallon of water. Allow all treated surfaces to air dry. Change solution daily or when solution becomes visibly dirty. For use with non-porous water resistant boots.

**To sanitize pre-cleaned and potable water-rinsed, non-porous public eating establishment and dairy food contact surfaces:** prepare a 200-400 ppm active quaternary solution by adding 1-2 ounces of this product to 4 gallons of water, 2.5-5 ounces to 10 gallons of water or 5-10 ounces to 20 gallons of water (or equivalent dilution).

**To prepare a 200, 300 or 400 ppm active quaternary solution,** use the following dilution table:

Active Quat Solution	1 Gallon	4 Gallons	10 Gallons	20 gallons
200 ppm	0.25 ounce	1 ounce	2.5 ounces	5 ounces
300 ppm	0.376 ounce	1.5 ounces	3.76 ounces	7.5 ounces
400 ppm	0.5 ounce	2 ounces	5 ounces	10 ounces

**To sanitize pre-cleaned mobile items in public eating establishments** (drinking glasses, dishes, eating utensils) immerse in a 200-400 ppm active quaternary solution for at least 60 seconds making sure to immerse completely. Remove items, drain the use-solution from the surface and air dry. Do not rinse.

**To sanitize pre-cleaned public eating establishment surfaces** (counters, tables, finished wood or plastic cutting boards), apply a 200-400 ppm active quaternary solution with a (cloth, sponge, low pressure coarse sprayer or hand pump trigger sprayer) making sure that the surface remains completely wet for at least 60 seconds and let air dry. Do not rinse.

**To sanitize pre-cleaned dairy processing equipment** (tanks, piping, pasteurizers, milking equipment) (apply a 200-400 ppm active quaternary solution with a low pressure coarse sprayer) or (circulate a 200-400 ppm active quaternary solution throughout the equipment). Surfaces must remain thoroughly wet for at least 60 seconds, followed by adequate draining and air drying. Do not rinse.

**Sanitization of beverage dispensing equipment:**

1. Wash equipment with a compatible detergent and rinse with potable water prior to sanitizing.
2. Fill equipment with a use-solution of 0.25–0.5 ounce of this product (or trade name) per gallon of water (or equivalent dilution).
3. Allow use-solution to remain in equipment for at least 60 seconds.
4. Drain thoroughly and allow to air dry before reuse. Do not rinse with potable water.

*<This statement is required for marketing food contact sanitizers in Wisconsin.>*

**WISCONSIN STATE DIVISION OF HEALTH**

**DIRECTIONS FOR EATING ESTABLISHMENTS**

1. Scrape and pre-wash utensils and glasses whenever possible.
2. Wash with a good detergent or compatible cleaner.
3. Rinse with clean water.
4. Sanitize in a solution of 1 ounce to 4 gallons of water (200 ppm active quaternary). Immerse all utensils for at least two minutes or for contact time specified by governing sanitary code.
5. Place sanitized utensils on a rack or drain board to air dry.
6. Fresh sanitizing solution should be prepared at least daily or when visibly soiled or diluted.

**NOTE:** A clean potable water rinse following sanitization is not permitted under Section HFS 196.13 of the Wisconsin Administrative Code.

**U. S. PUBLIC HEALTH SERVICE FOOD SERVICE SANITIZATION RECOMMENDATIONS**

**Cleaning and sanitizing:** Equipment and utensils must be thoroughly pre-flushed or pre-scraped and when necessary, pre-soaked to remove gross food particles and soil.

1. Thoroughly wash equipment and utensils in hot detergent solution.
2. Rinse utensils and equipment thoroughly with potable water.
3. Sanitize equipment and utensils by immersion in a use-solution of 1-2 ounces of this product per 4 gallons of water (or equivalent dilution) for at least 60 seconds at a temperature of 75°F.
4. For equipment and utensils too large to sanitize by immersion, apply a use-solution of 1-2 ounces of this product per 4 gallons of water (or equivalent dilution) by rinsing, spraying or swabbing until thoroughly wet.
5. Allow sanitized surface to drain and air dry. Do not rinse.

Bracketed information is denoted as: { insert as noted }, ( optional text ), [ required qualifier ].  
 The term "this product" used throughout this document may be replaced with the marketed product brand name.  
 The term "x" is a place holder for numerical digits.  
 Punctuation and plural/singular word forms may be adjusted to allow for grammatical correctness.  
 Appropriate metric or imperial unit conversion may be added as optional supplemental information.

**Dairy cow sanitizer application directions:** The udders, flanks and teats of dairy cows can be washed with a solution of 1 ounce of (this product or product name) in 4 gallons of warm water. Use a fresh towel for each cow. Avoid contamination of wash solution by dirt and soil. Do not dip used towel back into wash solution. When solution becomes visibly dirty, discard and provide fresh solution.

**To sanitize previously cleaned food-grade eggs** in shell egg and egg product-processing plants, spray with a solution of 1-2 ounces of (this product or product name) in 4 gallons of warm water (200-400 ppm quat). The solution must be warmer than the eggs, but not to exceed 130°F. Wet eggs thoroughly and allow to drain. Eggs sanitized with this product must be subjected to a potable water rinse only if they are broken immediately for use in the manufacture of egg products. Eggs must be reasonably dry before casing or breaking. The solution must not be reused for sanitizing eggs.

**To sanitize pre-cleaned immobile food processing equipment and surfaces** (tanks, finished wood or plastic chopping blocks, counter tops, conveyors) flood the area with a 200 to 400 ppm active quaternary solution for at least 60 seconds, making sure to wet all surfaces completely. Drain the use-solution from the surface and air dry.

**To sanitize pre-cleaned mobile food processing utensils** (knives, grinders, shredders, cleavers, ladles), immerse in a 200 to 400 ppm active quaternary solution for at least 60 seconds, making sure to immerse completely. Remove, drain the use-solution from the surface and air dry.

**Sanitization of food processing equipment, utensils, and other food contact articles regulated under 40CFR180.940:**

1. Scrape, flush, or pre-soak articles (whether mobile or stationary) to remove gross food particles and soil.
2. Thoroughly wash articles with a compatible detergent or cleaner.
3. Rinse articles thoroughly with potable water.
4. Sanitize by immersing articles with a use-solution of 1-2 ounces of this product (or trade name) per 4 gallons of water (or equivalent dilution) (200–400 ppm active quaternary) for at least 60 seconds. Articles too large for immersion should be thoroughly wetted by rinsing, spraying or swabbing.
5. Remove immersed items from solution to drain and then let air dry. Non-immersed items must be allowed to air dry.

**Sanitization of pre-cleaned sanitary filling equipment:** Prepare a use-solution of 0.25–0.5 ounce of this product per gallon of water (or equivalent dilution) (200-400 ppm active quaternary) for final washer and rinse applications. Allow surfaces to remain wet for at least 60 seconds. Drain thoroughly and allow to air dry before reuse. Do not rinse.

**Sanitization of pre-cleaned beer fermentation and storage tanks:** Recommended for sanitization of beer fermentation and holding tanks, citrus and food processing storage and holding tanks. Prepare a use-solution of 0.25–0.5 ounce of this product per gallon of water (or equivalent dilution) (200-400 ppm active quaternary) for mechanical or automated systems. Allow surfaces to remain wet for at least 60 seconds. Drain thoroughly and allow to air dry before reuse. No rinse is allowed. For mechanical operations or automated systems, the used sanitizing solution may not be reused for sanitization.

**Sanitization of food processing equipment:** Equipment and utensils must be thoroughly pre-flushed or pre-scraped and when necessary pre-soaked to remove gross food particles. Clean and rinse equipment thoroughly, then apply sanitizing solution containing 1-2 ounces of this product per 4 gallons of water (or equivalent use dilution) (200-400 ppm active quaternary). Allow surfaces to remain wet for at least 60 seconds. No rinse is allowed.

**For continuous treatment of meat and poultry or fruit and vegetable conveyors:** Remove gross food particles and excess soil by a pre-flush or pre-scrape, wash with a compatible detergent or cleaner. Rinse equipment thoroughly with potable water. Apply 1-2 ounces of this product per 4 gallons of water (or equivalent use dilution) (200-400 ppm active quaternary) to conveyors with suitable feeding equipment. Do not allow this solution to be sprayed directly on food. Controlled volumes of the sanitizer solution are applied to return portion of conveyor through nozzles so located as to permit maximum drainage of sanitizer solution from equipment and to prevent puddles on top of belt. During interruptions in operation, use a low pressure coarse sprayer to apply a use-solution of 1-2 ounces of this product per 4 gallons of water (or equivalent) (200-400 ppm active quaternary) to equipment, peelers, collators, slicers, and saws. Conveyor equipment should be free of product when applying this coarse spray. Allow surfaces to remain wet for at least 60 seconds. No rinse is allowed.

**Waterproof Glove Sanitizing Directions:** To prevent introducing harmful organisms into processing areas of food plants, waterproof gloves should be sanitized prior to entering or re-entering those areas. Remove gross contamination from gloves before sanitizing; then place gloved hand in a use-solution of 1 to 2 oz. of this product per 4 gallons (or equivalent dilution) (200-400 ppm quat) for 60 seconds (1 minute). Change the use-solution in the bath at least daily or more often if the solution appears visibly diluted or soiled.

### **Sanitization of Interior Hard, Non-Porous Surfaces of Water Softeners and Reverse Osmosis (RO) Units:**

**Water Softeners:** Sanitization should occur after initial installation, after the system is serviced and periodically during its use.

1. Backwash the softener and add 2 ounces of sanitizer per 8 gallons of water (200 ppm active quaternary) to the brine tank well. (NOTE: standard system capacity is 48 gallons). (The brine tank should have water in it to permit the solution to be carried into the softener).
2. Units/tanks must be washed with a compatible detergent and rinsed with potable water before sanitizing.
3. Proceed with the normal regeneration or interrupt the cycle after the brining step and let the softener soak for a minimum of 60 seconds.
4. Drain use-solution (sanitizing solution) from the unit.

**Reverse Osmosis (RO) Units:** Sanitization should occur after initial installation, after the system is serviced and periodically during its use.

1. Turn off RO system, drain storage tank and remove membrane element and pre-filters. Put membrane element in a plastic bag so it remains wet. Do not use this product to sanitize the membrane unit; it must be sterilized.
2. Units/tanks must be washed with a compatible detergent and rinsed with potable water before sanitizing.
3. Fill empty pre-filter housing with 2 ounces of sanitizer per 8 gallons of water (200 ppm active sanitizer) and turn on raw water. (NOTE: standard system capacity is 1-2 gallons).
4. After holding tank is full, let system stand idle for a minimum of 60 seconds. Turn off water. Drain holding tank.
5. Follow the manufacturer's directions for re-installation of new pre-filters, membrane element and post filter.

### **Sanitization of Interior Hard, Non-Porous Surfaces of Ice Machines, Water Coolers, Water Holding Tanks and Pressure Tanks**

**Ice Machines:** Sanitization should occur after initial installation, after the machine is serviced and periodically during its use.

1. Shut off incoming water line to machine.
2. Units/tanks must be washed with a compatible detergent and rinsed with potable water before sanitizing.
3. Prepare a solution of 2 ounces of sanitizer per 8 gallons of water (200 ppm active quaternary).
4. Circulate use-solution through machine making sure to wet all surfaces and allow a minimum contact of 60 seconds.
5. Drain the unit.

6. Return machine to normal operation.

**Water Coolers, Water Holding Tanks and Pressure Tanks:** Sanitization should occur after initial installation, after the system is serviced and periodically during its use.

1. Shut off incoming water line.
2. Units/tanks must be washed with a compatible detergent and rinsed with potable water before sanitizing.
3. Prepare a solution of 2 ounces of sanitizer per 8 gallons of water (200 ppm active quaternary).
4. Apply or circulate (if possible) to wet all surfaces and allow a minimum contact of 60 seconds.
5. Drain the unit.
6. Return to service by opening incoming water lines.

**<Not for Use in California>**

**Directions for fogging dairies, beverage and food processing plants:** Prior to fogging, food products and packaging material must be removed from the room or carefully protected. After cleaning, fog desired areas using 1 quart per 1000 cubic feet of room area with a solution containing 1.5 ounces of product to 1 gallon of water (1160 ppm active quaternary). Vacate the area of all personnel for a minimum of 2 hours after fogging. All food contact surfaces must be thoroughly rinsed with potable water before reuse.

**NOTE:** The fog generated is irritating to the eyes, skin and mucous membranes. Under no circumstances should a room or building be entered by anyone within two hours of the actual fogging. If the building must be entered, then the individuals entering the building must wear a self-contained respirator approved by NIOSH/MSHA, goggles, long sleeves and long pants.

**ALL SURFACES MUST BE CLEANED IN ACCORDANCE WITH LABEL DIRECTIONS PRIOR TO FOGGING. FOGGING IS AN ADJUNCT OR SUPPLEMENT TO NORMAL CLEANING PROCEDURES AND PRACTICES.**

**Mushroom Farm Premise Use:**

**Site Preparation:** The first step in any on-going sanitation program should be the removal of gross contamination and debris. This may be accomplished by using a shovel, broom, or vacuum, depending on the area to be disinfected.

**Cleaning and Disinfection:** For general disinfection, use 3 ounces of this product per 5 gallons of water (or equivalent dilution). Apply use-solution with a cloth, mop, sponge, lowpressure coarse sprayer or by immersion to thoroughly wet all surfaces. Treated surfaces must remain wet for a minimum of 10 minutes. Wipe or allow to air dry. For heavily soiled areas, pre-clean first. Prepare a fresh solution for each use or more often if use-solution becomes visibly diluted, clouded or soiled.

**For Heavy Duty Cleaning:** When greater cleaning is desired, use 6 ounces of this product per 5 gallons of water (or equivalent dilution). Heavily soiled areas may require repeated cleaning before treatment.

**DO NOT APPLY THE USE-SOLUTION TO THE MUSHROOM CROP, COMPOST OR CASING.**

Thoroughly rinse treated surfaces with potable water before they contact crop, compost or casing.

**Entryway Sanitizing Systems:** To prevent cross contamination from area to area, set the system to deliver (0.51 to 1.02 ounces per gallon of water) (or equivalent use dilution) (400 to 800 ppm active quaternary) of sanitizing solution. The (spray) (foam) should cover the entire path of the doorway. Set the system so that a continuous wet blanket of sanitizer solution is delivered to the floor. Do not mix other additives to the sanitizing solution.

**Citrus Canker Control:** For prevention of Citrus Canker Disease through treatment of equipment. This product, when used at a 1:48 dilution (2000 ppm active quaternary) or 2.67 ounces (79 ml) per gallon of water up to 200 ppm hardness (as CaCO<sub>3</sub>), is effective against *Xanthomonas axonopodis* pathovar *citri* (Citrus Canker). Treat all trucks, vehicles, and equipment thoroughly at the 1:48 dilution. Treatment can be applied to trucks, attached trailers, field harvesting equipment; including cargo

area, wheels, tires, under carriage, hood, roof, fenders and any other part of transportation equipment that can be taken into infested areas. To prevent the spread of Citrus Canker by this artificial means of transportation, treatments are made by trigger spraying, dipping or brushing, using the maximum rate of 2000 ppm quaternary. After use, all surfaces which come in contact with food or crop must be rinsed with potable water.

**Water And Smoke Damage Restoration:** For home, institutional, industrial and hospital use. This product is suitable for use in water damage restoration as a deodorizer on the following porous and semi-porous materials: carpets, cushions, sub floors, drywall, trim, frame lumber, tackless strip and paneling. Use a solution of 3 ounces of this product (or trade name) per 5 gallons of water (or equivalent dilution). Saturate all affected materials with enough solution to remain wet for a minimum of 10 minutes. Use proper ventilation.

**Smoke Damage Restoration:** Effective as a deodorizer for home, industrial, institutional and hospital use. This product is for use in smoke damage restoration situations to deodorize the following porous and semi-porous materials: carpets, carpet cushion, sub floors, drywall, trim and frame lumber, tackless strip and paneling. Follow directions in the Water Damage Restoration section. Using the recommended use-solutions, saturate affected materials with enough product to remain wet for a minimum of 10 minutes. Use proper ventilation.

**Sewer Backup And River Flooding:** During mitigation procedures, dilute 1.1 to 2.2 ounces of this product per gallon of water. Remove gross filth or heavy soil along with non-salvageable materials. Saturate all affected areas with the use-solution of (this product or product name). Apply the use-solution with a low pressure, coarse sprayer before and after cleaning and extraction.

**Carpet, Carpet Cushions And Other Porous Materials Such As Sub Floors, Drywall, Trim And Frame Lumber, Tackless Strip And Paneling:** For water damage from a clean water source, extract excess water. Test a hidden area for colorfastness. Remove gross filth or heavy soil. Dilute 1.1 to 2.2 ounces of this product per gallon of water (859-1718 ppm active quaternary). Apply the use-solution with a low pressure, coarse sprayer to fully saturate affected materials. Roll, brush or agitate into materials and allow the materials to remain damp for 10 minutes. Follow with a thorough extraction. Thoroughly dry all materials rapidly.

**Special Instructions For Carpet Deodorization:** This product may be used in homes, motels, hotels, nursing homes, schools and hospitals. This product is for use on wet, cleanable, synthetic fibers. Do not use on wool. Vacuum carpet thoroughly prior to cleaning. Test fabric for colorfastness.

- **For portable extraction units:** Mix 1.1 ounces per gallon of water.
- **For truck mounted extraction machines:** Mix 13.5 ounces per gallon of water and meter at 4 gallons per hour.
- **For rotary floor machines:** Mix 2.2 ounces per gallon of water and apply at the rate of 300-500 sq. ft. per gallon.

Do not mix (this product or product name) with other cleaning products. Follow the cleaning procedures specified by the manufacturer of the cleaning equipment. After using this product set the carpet pile in one direction with a stiff brush and protect the carpet from furniture legs and bases while drying. Do not over wet. If applied to stain resistant nylon carpet, apply fabric protector according to the carpet manufacturer's directions.

**Carpet Deodorization for Home, Industrial and Institutional Use and Hospital Use:** This product may be used in homes, motels, hotels, nursing homes and hospitals.

Vacuum carpet thoroughly prior to application. Mix 1.1 ounces per gallon of water. Follow injection and/or extraction procedures as specified for any conventional steam cleaning equipment you are using. For rotary floor machines, mix 2.2 ounces per gallon of water and spray on carpet at a rate of 300-500 sq. ft. per gallon.

For use on washable synthetic fibers. Do not use on wool. Test colorfastness of carpet before use. Apply diluted product to a small concealed spot, then rub with a clean white cloth. If color changes or transfers to cloth, a water based product should not be used.

After using the product, set carpet pile in one direction with a stiff brush. Place aluminum foil under the legs of furniture while carpet is drying. Over-wetting can cause carpet to shrink. Manufacturer assumes no responsibility for over-wetting misuse.

**NOTE:** This product should not be mixed with other cleaning chemicals.

**Hatchery Rooms Using Fogging Devices:** Remove all animals and feed from premises, vehicles and enclosures. Remove all litter and manure from floors, walls and surfaces of the room to be treated. Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean all surfaces with a compatible cleaner or detergent and rinse thoroughly with potable water. Close off room so fog is confined to room to be treated. Mix 140 ounces of (this product or product name) to 2.5 gallons of water. Insert the fogger nozzle through a suitable opening into the room. With the setting in maximum output, fog for one minute for each 4000 cubic feet of room space. When fogging is completed, ventilate buildings and other closed spaces. Do not house livestock or employ equipment until treatment has been absorbed or dried. Thoroughly clean all surfaces with a compatible soap or detergent and rinse thoroughly with potable water.

**Incubators and Hatcheries Using Fogging Devices:** Mix 13.5 ounces of (this product or product name) to 120 ounces of water. Fog 3 to 8 ounces of this use-solution into setters and hatchers immediately after transfer. Repeat daily in setters and every 12 hours in hatchers. Discontinue hatcher treatments at least 24 hours prior to pulling the hatch.

It is acceptable to fog hatchers and setters with a 1.1 ounces per gallon use-solution of (this product or product name) on an hourly or every other hour basis. If this is done, fog for 30-90 seconds once per hour or once every two hours. When fogging is complete, ventilate buildings, and other closed spaces. Do not house livestock or employ equipment until treatment has been absorbed or dried. Thoroughly scrub all treated feed racks, mangers, troughs, automatic feeders, fountains and waterers with a compatible soap or detergent, and thoroughly rinse with potable water. Best results are achieved with use-solution water temperature ranging from 78-110°F.

**NOTE:** The fog generated is irritating to the eyes, skin and mucous membranes. Under no circumstances should a room or building be entered by anyone within two hours of the actual fogging. If the building must be entered, the individuals entering the building must wear a self-contained respirator approved by NIOSH/MSHA, goggles, long sleeves and long pants.

**ALL SURFACES MUST BE CLEANED IN ACCORDANCE WITH LABEL DIRECTIONS PRIOR TO FOGGING. FOGGING IS AN ADJUNCT OR SUPPLEMENT TO NORMAL PROCEDURES AND PRACTICES.**

## **DILUTION SYSTEM SPECIFIC DIRECTIONS FOR USE**

**PACKETS:** *Pre-measured tear open packets* – (Simply) (tear) (open and) pour contents (of packet) into x gallons of water (for a (200ppm)(xppm)(547ppm) dilution). Keep packets in box until ready to use.

### **INTERMEDIATE SERVICE SOLUTION CONTAINER:**

1. Create Intermediate Service Solution by combining (x oz product) (x packet(s)) and x (oz) (gal) water for a (x:x) intermediate dilution.
2. Follow Directions for Use for (Sanitization), (General Disinfection), (Hospital Disinfection), (Insert applicable use header).

	Minimum Level	Intermediate Level	Maximum Level
Sanitization	200ppm	xppm	400ppm
General Disinfection	467ppm	xppm	546ppm
	Minimum Level		
Hospital Disinfection	547 ppm	xppm (with x greater than 547 ppm)	

3. Combine x oz of the Intermediate Service Solution and x (oz)(gal) water to achieve the final use active concentration of (200ppm) (xppm) (547ppm)

## **PRECAUTIONARY STATEMENTS**

### **PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS.**

#### **DANGER**

KEEP OUT OF REACH OF CHILDREN. CORROSIVE. Causes irreversible eye damage and skin burns. Do not get in eyes, on skin or on clothing. Wear goggles or face shield, rubber gloves, and protective clothing. Harmful if swallowed. Remove contaminated clothing and wash before reuse. Wash thoroughly with soap and water after handling.

*<Required for products supplied in containers of 5 gallons or greater>*

#### **ENVIRONMENTAL HAZARDS**

This product is toxic to fish. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

Contains no phosphorus.

#### **STORAGE AND DISPOSAL**

**DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL.**

**PESTICIDE STORAGE** - Store in a dry place no lower in temperature than 50°F or higher than 120°F. Store in a tightly closed container in an area inaccessible to children.

**PESTICIDE DISPOSAL** - Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

#### **CONTAINER DISPOSAL - <mandatory language - choose applicable directions>**

<i>For products only with household/residential use sites</i>	Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available.
<i>For products with industrial, institutional, and commercial use sites</i>	Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse container promptly after emptying.  For containers 2 gallons or less with a one-way opening that doesn't allow containers to be rinsed: Nonrefillable container. Do not reuse or refill this container. Replace cap and discard in trash. Offer for recycling if available.
<i>For containers 5 gallons or less</i>	Triple rinse as follows: Fill container ¼ full with water and recap. Agitate vigorously. Follow Pesticide Disposal instruction for rinsate disposal. Drain for 10 seconds after the flow begins to drip. Repeat procedure two more times. Offer for recycling or reconditioning, if available. If not available, puncture and dispose in a sanitary landfill.
<i>For containers greater than 5 gallons</i>	Triple rinse as follows: Fill container ¼ full with water. Tip container on its side and roll it back and forth ensuring at least one complete revolution for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Follow Pesticide Disposal instructions for rinsate disposal. Repeat this procedure two more times. Offer for recycling or reconditioning, if available. If not available, puncture and dispose in a sanitary landfill.

*<NOTE: spill or leak procedures is required language>*

#### **SPILL OR LEAK PROCEDURES**

Small spills may be mopped up, flushed away with water or absorbed on some absorbent material and incinerated. Large spills should be contained, the material then moved into containers and disposed of by approved methods for hazardous wastes.

<when applicable>

**PACKET DISPOSAL** – Do not reuse; discard empty container and put in trash.

## **GRAPHICS**

<NOTE TO REVIEWER: Product label may show graphic depicting the surface/site application appropriate with direction for use, PPE or package type. Graphics may include but are not limited to the depictions below.>

(kitchen)  
(bathroom)  
(shower curtain)  
(floor)  
(wall)  
(stainless steel surfaces)  
(glazed tiles)  
(glazed  
porcelain)  
(plastic surfaces)  
(bathtubs)  
(cabinets)  
(tables)  
(chairs)  
(packet)  
(packet ripped open)  
(packet being poured)  
(intermediate service solution container)  
(spray dispensing container)  
(dispensing of product from intermediate service solution container to spray dispensing container)  
(sink with faucet)  
(tub)



<Note to reviewer: The following may be used only if the supplemental registrant has obtained a Kosher certification.>

<Any Kosher Symbol such as:>



**PACKET LABEL UNIT PACKAGING**

**(1)(X) OUNCE PACKET LABEL TO BE USED WITH MASTER CONTAINER LABEL:  
MIX EACH PACKET WITH (4)(X) GALLONS OF WATER:**

*(Note to reviewer: x ounce packets and corresponding x amount of water will be adjusted to yield equivalent dilution in accordance with directions for use on this label)*

**BTC® 2125M 10% Solution  
1839-86**

**ACTIVE INGREDIENTS**

Alkyl (60% C <sub>14</sub> , 30% C <sub>16</sub> , 5% C <sub>12</sub> , 5% C <sub>18</sub> ) dimethyl benzyl ammonium chlorides.....	5.0%
Alkyl (68% C <sub>12</sub> , 32% C <sub>14</sub> ) dimethyl ethylbenzyl ammonium chlorides.....	5.0%
<b>(INERT)(OTHER) INGREDIENTS</b> .....	<u>90.0%</u>
<b>TOTAL</b> .....	<b>100.0%</b>

**KEEP OUT OF REACH OF CHILDREN  
DANGER**

*(Note to reviewer: Only one of the following statements is required. Referral to "Directions for Use" is required when directions for use are only found in booklet. Choose applicable statement noted in curly bracket.)*

See (side) (back) panel for additional precautionary statements.

or

See booklet for {(complete) (additional)} information (and) (including) (Directions for Use)

**Disposal** – Do not reuse; discard empty packet by putting in trash.

EPA REG. NO. 1839-86

EPA EST. NO. (1839-IL-01)(xxxx-xx-xxx)

STEPAN COMPANY  
22 West Frontage Road • Northfield, IL 60093  
(847) 446-7500

NOT FOR INDIVIDUAL SALE

Pour contents of packet into xx (gallons)(gal)(oz) of water for a (x) dilution

NET CONTENTS: xx (ounces)(oz) per packet

Bracketed information is denoted as: { insert as noted }, ( optional text ), [ required qualifier ].  
The term "this product" used throughout this document may be replaced with the marketed product brand name.  
The term "x" is a place holder for numerical digits.  
Punctuation and plural/singular word forms may be adjusted to allow for grammatical correctness.  
Appropriate metric or imperial unit conversion may be added as optional supplemental information.